

Multi-Index Quasi-Monte Carlo

An algorithm for simulating PDEs with random coefficients

Pieterjan Robbe

joint work with D. Nuyens and S. Vandewalle

✉ pieterjan.robbe@cs.kuleuven.be



August 16, 2016



The missing link?

	Monte Carlo	Quasi-Monte Carlo
Multilevel	MLMC ^{a,b}	MLQMC ^{c,d,e}
Multi-index	MIMC ^f	?

- ^a K. A. Cliffe, M. B. Giles, R. Scheichl and A. L. Teckentrup, *Multilevel Monte Carlo Methods and Applications to Elliptic PDEs with Random Coefficients*, Computing and Visualization in Science, 14 (2011), pp. 3-15. [Cliffe, 2011]
- ^b A. Barth, C. Schwab and N. Zollinger, *Multilevel Monte Carlo Finite Element Method for Elliptic PDEs with Stochastic Coefficients*, Numerische Mathematik, 119 (2011), pp. 123-161. [Barth, 2011]
- ^c F. Y. Kuo, R. Scheichl, C. Schwab, I. H. Sloan and E. Ullmann, *Multilevel Quasi-Monte Carlo Methods for Lognormal Diffusion Problems*, Mathematics of Computation, under revision, 2016. [Kuo, 2016]
- ^d F. Y. Kuo and D. Nuyens, *Application of Quasi-Monte Carlo Methods to Elliptic PDEs with Random Diffusion Coefficients – A Survey of Analysis and Implementation*, Foundations of Computational Mathematics, to appear, 2016. [Kuo, 2016/2]
- ^e I. G. Graham, F. Y. Kuo, J. A. Nichols, R. Scheichl, C. Schwab and I. H. Sloan, *Quasi-Monte Carlo Finite Element Methods for Elliptic PDEs with Lognormal Random Coefficients*, Numerische Mathematik, 131 (2014), pp. 329-368. [Graham, 2014]
- ^f A.-L. Haji-Ali, F. Nobile and R. Tempone, *Multi-Index Monte Carlo: When Sparsity Meets Sampling*, Numerische Mathematik 132 (2016), pp. 767-806. [Haji-Ali, 2016]

Contents

1. PDEs with Random Coefficients

An example from geophysics

2. From MLMC to MIMC

Exploiting coarser grids for variance reduction

3. From MIMC to MIQMC

Improving the convergence rate

4. Results

Numerical results

5. Conclusions

Some closing thoughts

PDEs with Random Coefficients

PDEs with random coefficients

- Let (Ω, \mathcal{A}, P) be a complete probability space and $D \subset \mathbb{R}^d$ a bounded domain
- We look for solutions $u : D \rightarrow \mathbb{R}$ that solve almost surely (a.s.)

$$-\nabla \cdot (a(\mathbf{x}, \omega) \nabla u(\mathbf{x}, \omega)) = f(\mathbf{x}) \quad \text{for } \mathbf{x} \in D \text{ and } \omega \in \Omega$$


with given boundary conditions

$$u(\mathbf{x}, \cdot) = u_1(\mathbf{x}) \quad \text{for } \mathbf{x} \in \partial_1 D$$

$$n(\mathbf{x}) \cdot (a(\mathbf{x}, \cdot) \nabla u(\mathbf{x}, \cdot)) = u_2(\mathbf{x}) \quad \text{for } \mathbf{x} \in \partial_2 D$$

- The **goal** is to compute statistics of the functional $G = \mathcal{G}(u)$ for some sufficiently smooth a and \mathcal{G}

Flow through porous media



- ❖ Represent permeability of the porous medium as a lognormal **random field** $k(\mathbf{x}, \omega) = \exp(Z(\mathbf{x}, \omega))$, where $Z(\mathbf{x}, \omega)$ is an underlying Gaussian random field
- ❖ A Gaussian random field is characterised by its **mean** $\mu(\mathbf{x}) = \mathbb{E}[Z(\mathbf{x})]$ and **covariance function** $\text{cov}(Z(\mathbf{x}), Z(\mathbf{y}))$
- ❖ Every fixed $\omega \in \Omega$ yields a deterministic **realisation** of the random field

On the choice of covariance function

- ❖ A common choice is the **exponential covariance** function

$$C(\rho) = \sigma^2 \exp\left(\frac{\rho}{\lambda}\right)$$

with $\rho = \|\mathbf{x} - \mathbf{y}\|_p$ and p the usual ℓ_p -norm

- ❖ We choose the more general **Matérn covariance** function


$$C(\rho) = \sigma^2 \frac{1}{2^{\nu-1} \Gamma(\nu)} \left(\sqrt{2\nu} \frac{\rho}{\lambda} \right)^{\nu} K_{\nu} \left(\sqrt{2\nu} \frac{\rho}{\lambda} \right)$$

with Γ the Gamma function and K_{ν} the modified Bessel function of the second kind


- ❖ The Matérn covariance reduces to the exponential case when $\nu = 1/2$, $\nu = \infty$ is the squared exponential case.

On the choice of covariance function


Some examples of $k(\mathbf{x}, \omega)$ for different parameter sets



$$\begin{aligned}\lambda &= 1 \\ \sigma^2 &= 1 \\ \nu &= 2.5\end{aligned}$$



$$\begin{aligned}\lambda &= 0.3 \\ \sigma^2 &= 1 \\ \nu &= 1\end{aligned}$$



$$\begin{aligned}\lambda &= 0.075 \\ \sigma^2 &= 1 \\ \nu &= 0.5\end{aligned}$$


The KL expansion

- Classical technique to take samples from $k(\mathbf{x}, \omega)$ is the Karhunen–Loève or **KL-expansion**

$$k(\mathbf{x}, \omega) = \bar{k} + \exp \left(\sum_{r=1}^{\infty} \sqrt{\theta_r} f_r(\mathbf{x}) \xi(\omega) \right)$$

with θ_r and f_r the solutions of the Fredholm equation

$$\int_D C(\mathbf{x}, \mathbf{y}) f_r(\mathbf{y}) d\mathbf{y} = \lambda_r f_r(\mathbf{y}), \quad \mathbf{x} \in D$$



$r = 1$

$r = 2$

$r = 4$


$r = 15$

$r = 88$

The KL expansion

- Approximation quality of the KL expansion determined by eigenvalue decay rate

$$\theta_r \sim \mathcal{O}\left(r^{-\frac{4\nu+d}{d}}\right)$$



Darcy flow

- Steady-state flow through porous media given by **Darcy's law**

$$-\nabla \cdot k(\mathbf{x}, \omega) \nabla p(\mathbf{x}, \omega) = f(\mathbf{x})$$

with k the hydraulic conductivity and p the pressure head

- This is an elliptic **PDE with random coefficients!**
- Let us choose $D = [0, 1]^3$ and the functionals

$$G^1 = p(\mathbf{x}^*), \quad \mathbf{x}^* = (0.5, 0.5, 0.5)$$

$$\text{and} \quad G^2 = - \int_0^1 \int_0^1 k \frac{\partial p}{\partial x} \Big|_{x=1} dy dz$$

- Our **goal** is to compute $\mathbb{E}[G^1]$ and $\mathbb{E}[G^2]$


From MLMC to MIMC

Numerical approximation


- ❖ We assume that different approximations of u are available based on the discretization parameter h
- ❖ For MLMC, one usually assumes a **geometrical** mesh hierarchy

$$h_\ell = h_0 2^{-\ell}$$


- ❖ The approximations are indexed by a **level** $\ell \in \mathbb{N}$



$\ell = 0$



$\ell = 1$



$\ell = 2$

Multilevel Monte Carlo

- Suppose we denote by G_ℓ the approximation of G at level ℓ
- Let $\Delta G_\ell := \begin{cases} G_\ell & \text{if } \ell = 0 \\ G_\ell - G_{\ell-1} & \text{if } \ell > 0 \end{cases}$
- Using the **telescoping sum** identity,

$$\mathbb{E}[G_L] = \sum_{\ell=0}^L \mathbb{E}[\Delta G_\ell]$$

the MLMC [Giles, 2008] estimator can be expressed as

$$\hat{G}_{\text{ML}} = \sum_{\ell=0}^L \frac{1}{N_\ell} \sum_{n=1}^{N_\ell} \Delta G_\ell(\omega_{\ell,n})$$


- Main point:** MLMC is a recursive control variate strategy that uses a hierarchy of coarser grids

Numerical approximation (2)


- ❖ We assume that different approximations of u are available based on the discretization parameters $h_i, i = 1 \dots d$
- ❖ Similar to MLMC, assume a **geometrical** mesh hierarchy

$$h_{\ell_i} = h_0 2^{-\ell_i}$$


- ❖ Approximations are indexed by a **multi-index** $\vec{\ell} = (\ell_i)_{i=1}^d \in \mathbb{N}$



$$\ell = (0, 0)$$




$$\ell = (1, 1)$$



$$\ell = (2, 2)$$

Multi-Index Monte Carlo



Multi-Index Monte Carlo

- Suppose we denote by $G_{\vec{\ell}}$ the approximation of G at index $\vec{\ell}$
- Let $\Delta G_{\vec{\ell}} := \left(\bigotimes_{i=1}^d \Delta_i \right) G_{\vec{\ell}}$ with

$$\Delta_i G_{\vec{\ell}} = \begin{cases} G_{\vec{\ell}} & \text{if } \ell_i = 0 \\ G_{\vec{\ell}} - G_{\vec{\ell} - \vec{e}_i} & \text{if } \ell_i > 0 \end{cases} \quad \text{for } i = 1 \dots d$$

- The MIMC estimator [Haji-Ali, 2016] can be expressed as

$$\hat{G}_{\text{MI}} = \sum_{\vec{\ell} \in \mathcal{I}} \frac{1}{N_{\vec{\ell}}} \sum_{n=1}^{N_{\vec{\ell}}} \Delta G_{\vec{\ell}}(\omega_{\vec{\ell},n})$$

where $\mathcal{I} \subset \mathbb{N}^d$ is an appropriate set of indices, the **index set**

Example

- For $d = 2$ and $\vec{\ell} = (1, 2)$, we have

$$\begin{aligned}\Delta G_{(1,2)}(\omega) &= \Delta_2 (\Delta_1 G_{(1,2)}(\omega)) \\ &= \Delta_2 (G_{(1,2)}(\omega) - G_{(0,2)}(\omega)) \\ &= (G_{(1,2)}(\omega) - G_{(0,2)}(\omega)) - (G_{(1,1)}(\omega) - G_{(0,1)}(\omega)) \\ &= G_{(1,2)}(\omega) - G_{(0,2)}(\omega) - G_{(1,1)}(\omega) + G_{(0,1)}(\omega)\end{aligned}$$

- Key point** is that these 4 solutions are based on the same realisation of the random field $k(\cdot, \omega)$
- In general, computing a single realisation of $\Delta G_{\vec{\ell}}$ requires the solution of 2^d different PDEs, where the cost is dominated by the solution at index $\vec{\ell}$

Complexity analysis

- Classical MSE error splitting: $\text{MSE} = \mathbb{V}[\hat{G}] + \mathbb{E}[(\hat{G} - G)]^2$
- By independence, we have $\mathbb{V}[\hat{G}] = \sum_{\vec{\ell} \in \mathcal{I}} \frac{V_{\vec{\ell}}}{N_{\vec{\ell}}}$ with $V_{\vec{\ell}} := \mathbb{V}[\Delta G_{\vec{\ell}}]$
- The second term can be approximated as


$$\mathbb{E}[(\hat{G} - G)] = \left| \sum_{\vec{\ell} \notin \mathcal{I}} \mathbb{E}[\Delta G_{\vec{\ell}}] \right| \approx \left| \sum_{\vec{\ell} \in \partial \mathcal{I}} \mathbb{E}[\Delta G_{\vec{\ell}}] \right|$$

with $\partial \mathcal{I}$ the **boundary** of the index set \mathcal{I} . Note: this requires sufficient decay of the $\mathbb{E}[\Delta G_{\vec{\ell}}]$ (e.g., exponential, see [Haji-Ali, 2016])


- Conclusion:** error and work complexity depends on choice of index set!

On the choice of index set


$$\mathcal{I}(L) = \left\{ \vec{\ell} \in \mathbb{N}^d : \ell_i \leq L \text{ for all } 1 \leq i \leq d \right\}$$



$$\mathcal{I}(L) = \left\{ \vec{\ell} \in \mathbb{N}^d : \sum_{i=1}^d \ell_i \leq L \right\}$$






$$\mathcal{I}(L) = \left\{ \vec{\ell} \in \mathbb{N}^d : \prod_{i=1}^d \max(1, \ell_i) \leq L \right\}$$



Boundary of an arbitrary index set

- ❖ \mathcal{I} is valid when it is **downward closed**
- ❖ An index $\vec{\ell}$ **dominates** an index $\vec{\tau}$ in a certain direction i if $\ell_i > \tau_i$
- ❖ An index $\vec{\ell}$ is an **interior index** if it is dominated by another index in the direction of the position of its maximum entry
- ❖ The **boundary** $\partial\mathcal{I}$ of an index set \mathcal{I} are these indices $\vec{\ell} \in \mathcal{I}$ that are part of the index set but are not interior indices



From MIMC to MIQMC

The MIQMC estimator

- Let us assume that we have a **randomly shifted rank-1 lattice rule** available that approximates

$$I_s(f) = \int_{[0,1]^s} f(\mathbf{y}) d\mathbf{y} \approx \frac{1}{K} \sum_{k=0}^{K-1} \frac{1}{N} \sum_{n=0}^{N-1} f(\{\mathbf{t}_n + \mathbf{\Delta}_k\})$$

(see tutorials on Sunday, or first plenary talk by D. Nuyens)

- In our setting, the “integral” is the expectation

$$\begin{aligned} \mathbb{E}[G_{\vec{\ell}}] &= \int_{\mathbb{R}^\infty} G_{\vec{\ell}}(u(\cdot, \xi_1, \xi_2 \dots)) d\Phi(\xi_1, \xi_2 \dots) \\ &\approx \int_{[0,1]^s} G_{\vec{\ell}}(u(\cdot, \Phi^{-1}(y_1), \dots, \Phi^{-1}(y_s), 0, \dots)) d\mathbf{y} \end{aligned}$$

with Φ the cumulative normal density and we apply the change of variables component-wise

$$\boldsymbol{\xi} = \Phi^{-1}(\mathbf{y}) = (\Phi^{-1}(y_1), \Phi^{-1}(y_2), \dots) \in \mathbb{R}^{\mathbb{N}} \text{ and } \mathbf{y} \in (0, 1)^{\mathbb{N}}$$

The MIQMC estimator


- ❖ The MIQMC estimator reads

$$\hat{G}_{\text{MI}}^* = \sum_{\vec{\ell} \in \mathcal{I}} \frac{1}{K} \sum_{k=0}^{K-1} \frac{1}{N_{\vec{\ell}}} \sum_{n=0}^{N_{\vec{\ell}}-1} \Delta G_{\vec{\ell}}(\Phi^{-1}(\{\mathbf{t}_n + \mathbf{\Delta}_{k,\vec{\ell}}\}))$$

- ❖ For linear functionals G in a single- or multilevel setting, it can be shown that the integrand belongs to a weighted Sobolev space with (S)POD-weights, see [Graham, 2014], [Kuo, 2016] and [Kuo, 2016/2]
- ❖ Software for generating these lattice rules: QMC4PDE
- ❖ The convergence rate depends on decay rate of eigenvalues, but is independent of the number of dimensions s


Results

Numerical results (1)




- ❑ Three-dimensional problem with Dirichlet boundary conditions
- ❑ Quantity of interest is G^1 (point evaluation)
- ❑ Lognormal random field with Matérn covariance, $\lambda = 1, \nu = 2.5, s = 12$
- ❑ Solve deterministic PDE using FV, CG with AMG preconditioner
- ❑ Coarsest mesh has $4^3 = 64$ DOF, finest mesh has $128^3 \approx 2\text{M}$ DOF

Numerical results (2)



- Three-dimensional problem with Dirichlet boundary conditions
- Quantity of interest is G^1 (point evaluation)
- Lognormal random field with Matérn covariance, $\lambda = 0.3, \nu = 1, s = 201$
- Solve deterministic PDE using FV, CG with AMG preconditioner
- Coarsest mesh has $4^3 = 64$ DOF, finest mesh has $128^3 \approx 2\text{M}$ DOF

Numerical results (3)



- Three-dimensional **flow cell** geometry
- Quantity of interest is G^2 (flux through boundary)
- Lognormal random field with Matérn covariance, $\lambda = \mathbf{0.075}$, $\sigma^2 = 1$, $\nu = \mathbf{0.5}$, $s = \mathbf{3000}$
- Solve deterministic PDE using FV, CG with AMG preconditioner
- Coarsest mesh has $4^3 = 64$ DOF, finest mesh has $128^3 \approx 2\text{M}$ DOF

Conclusions

Closing Thoughts

- ❖ MIQMC is a QMC extension of MIMC [Haji-Ali, 2016] or a MIMC extension of MLQMC [Kuo, 2016]
- ❖ We show numerical evidence that the MIQMC estimator achieves a convergence rate as good as MIMC, and as good as (or better than) MLQMC for smooth problems
- ❖ Cost and error analysis of the estimator must take into account the shape of the index set (can be complicated)
- ❖ A general-purpose library is available on <https://github.com/PieterjanRobbe/MultilevelEstimators.jl>
preprint is available on ArXiv
<http://arxiv.org/abs/1608.03157>
- ❖ Future work: adaptive construction of the index set \mathcal{I}